Supplementary Material to "Association of TYK2 polymorphisms with autoimmune diseases: A comprehensive and updated systematic review with meta-analysis"

Table S2 - Newcastle-Ottawa and Clark-Baudouin quality assessment scale for the studies included in the meta-analysis.

Author	Year	Newcastle-Ottawa Score	Clark-Baudouin Score
Almlöf et al.	2017	9	10
Alonso-Pérez et al.	2014	8	9
Ban et al.	2009	8	9
Can et al.	2015	8	9
Contreras-Cubas et al.	2019	9	10
Couturier et al.	2011	8	9
Diogo et al.	2015	9	10
Ellinghaus et al.	2016	9	10
Graciolo et al.	2019	8	8
Graham et al.	2011	7	8
Harley et al.	2008	9	9
Hellquist et al.	2009	8	9
Järvinen et al.	2010	7	8
Kyogoku et al.	2009	8	8
Langefeld et al.	2017	9	10
Li et al.	2011	7	8
Lian et al.	2013	6	7
López-Isac et al.	2016	8	9

Author	Year	Newcastle-Ottawa Score	Clark-Baudouin Score
Mero et al.	2010	8	8
Mohamadhosseini et al.	2019	9	9
Myrthianou et al.	2017	8	9
Nagafuchi et al.	2015	8	8
Prieto-Pérez et al.	2015	9	10
Qiu et al.	2013	9	10
Sato et al.	2009	8	9
Shaiq et al.	2013	7	8
Sigurdsson et al.	2005	8	9
Sigurdsson et al.	2007	8	9
Strange et al.	2010	9	9
Suarez-Gestal et al.	2009	8	9
Suarez-Gestal et al.	2009	9	10
Tang et al.	2015	9	9
Westra et al.	2018	9	10
Zaplakhova <i>et al</i> .	2018	8	9

Quality of articles were calculated using the Newcastle-Ottawa Scale (NOS), as described in the Methods Section. Selection: minimum 1 – maximum 4 stars; Comparability: 0-1 star; and Exposure: 1-3 stars. In addition, the Clark-Baudouin Score (CBS) was also used to assess the quality of the studies. The CBS consists in a scoring systems that allows from 0 to 10 points regarding pre-defined criteria.